


Improving Pollination Quality in Seed Crops



Riley Reed

Beeinformed.org

About me



A colony is alive, a hive is a box.



Swarming



Frames



Supers



Pollen trap



Buzzingacrossamerica.com



beeequipmentsaustralia.com

Supplemental feed



teeth.org.au/sugar



Breakdown

- Part one: Supplemental feeding
 - Chapter 1: Supplemental feeding in carrot seed
 - Chapter 2: Supplemental feeding in canola
 - Chapter 3: Supplemental feeding and colony health
- Part two: Hive space requirements
 - Chapter 4: Supering during carrot seed

Part 1: Supplemental Feeding

Why should I care?

- ~1/3 of crops depend on or at least benefit from pollinators (Klein et al, 2007)
- Honey bees are the most widely used pollinator
- US vegetable seed is worth over \$125,000,000 (USDA NASS 2019)
- US carrots are worth over \$750,000,000 (USDA NASS 2019)




Pollen transfer between fields lowers seed purity.



Benedict Vanheems
growveg.com

almanac.com/plant/carrots

Transgenic Crops



The screenshot shows a web browser window with the URL agriculture.basf.us/crop-protection/Campaigns/canola-stewardship/canola-stewardship-media-statements/canola-stewardship-upd. The page features the BASF logo and tagline "We create chemistry" at the top. A navigation menu includes "Crops", "Products", "Agronomics", "About BASF Ag Solutions", "Product Labels", and "AgSolutions Finder". The main heading is "Omega-3 events (LBFLFK and NS-B50027-4) in BASF Canola Seed". The text below the heading states: "RESEARCH TRIANGLE PARK, NC, April 13, 2021 – BASF has detected the presence of 2 distinct events (more commonly referred to as GMO traits) in canola seed samples taken from BASF's hybrid seed production fields in the United States. These events are designed to increase Omega-3 fatty acid content in the seed oil. While the detected events, LBFLFK or BASF-Cargill Omega-3 and NS-B50027-4, owned by Nuseed, are approved for cultivation in the United States, they have not yet received approval in key export countries." It further notes that "All affected 2020 hybrid seed lots were isolated and contained on farm or recalled from the commercial channel. BASF is working with customers to replace this recalled seed in time for the 2021 planting season." and "Based on these findings and information regarding additional field activities with the Omega-3 events in 2018 and 2019, we immediately initiated testing of retained lots from hybrid canola seed produced in these years in Montana and Washington State." It also mentions that "Until February 2021, no Nuseed authorized third-party laboratory was available to test for Nuseed's NS-B50027-4 event. BASF's investigation and retained seed sample testing is ongoing." The bottom of the page shows the start of a sentence: "BASF is working with the canola seed distribution channel on product recalls in an effort to prevent seed".

Omega-3 events (LBFLFK and NS-B50027-4) in BASF Canola Seed

RESEARCH TRIANGLE PARK, NC, April 13, 2021 – BASF has detected the presence of 2 distinct events (more commonly referred to as GMO traits) in canola seed samples taken from BASF's hybrid seed production fields in the United States. These events are designed to increase Omega-3 fatty acid content in the seed oil. While the detected events, LBFLFK or BASF-Cargill Omega-3 and NS-B50027-4, owned by Nuseed, are approved for cultivation in the United States, they have not yet received approval in key export countries.

All affected 2020 hybrid seed lots were isolated and contained on farm or recalled from the commercial channel. BASF is working with customers to replace this recalled seed in time for the 2021 planting season.

Based on these findings and information regarding additional field activities with the Omega-3 events in 2018 and 2019, we immediately initiated testing of retained lots from hybrid canola seed produced in these years in Montana and Washington State.

Until February 2021, no Nuseed authorized third-party laboratory was available to test for Nuseed's NS-B50027-4 event. BASF's investigation and retained seed sample testing is ongoing.


BASF is working with the canola seed distribution channel on product recalls in an effort to prevent seed



Honey bees can fly up to 9 kilometers.

● Optimal Foraging Theory
2-3km


● Geometric Framework
Up to 9.5km




Monocultures and hybrid production decrease food resources.




Poor nutrition is not immediately evident, but the effects can be long lasting




The 4 P's



Feeding sugar syrup increased the collection of kiwi pollen.



Feeding pollen substitute decreased the collection of other pollen other plants.



Going forward, two different cropping systems will be used.



Goals

- Improved isolation between fields
- Decreased risk of transgene escape
- Healthier, more valuable, colonies




Chapter 1: Carrot seed



Hypothesis 1 : Providing supplemental in-hive feeding will increase the number of foragers working in the field without impacting overall foraging activity.






 Fed


 Unfed

 Field

How does feeding impact foraging behavior?



The waggle dance is used to communicate the location and quality of a resource




Waggle Dance Game:

<https://askbiologist.asu.edu/bee-dance-game/how-to-play.html>



Observation hives allow to observe the inner workings of a hive



Waggle dances can also be used to decode foraging behavior




Expected results



Hypothesis 2 : Providing supplemental in-hive feeding will decrease the amount of non-carrot pollen collected.



How does feeding impact pollen collection?



Kosher salt
of table salt.

Nutrition Facts	
1133 servings per container	
Serving size	1/4 tsp (1.2g)
Amount per serving	
Calories	0
% Daily Value*	
Total Fat 0g	0%
Sodium 480mg	21%
Total Carbohydrate 0g	0%
Protein 0g	


Not a significant source of saturated fat, trans fat, cholesterol, dietary fiber, total sugars, added sugars, vitamin D, calcium, iron and potassium.

*The % Daily Value (DV) tells you how much a nutrient in a serving of food contributes to a daily diet. 2,000 calories a day is used for general nutrition advice.

INGREDIENTS: SALT, YELLOW PRUSSIAN OF SODA (AN ANTICAKING AGENT).

MORTON SALT, INC. CHICAGO, IL 60606
QUESTIONS CALL 1-800-775-7753


Expected results



Summer 2022



Colored powders were removed too quickly.



Månefjord et al. 2022



Hand painting requires too much time to mark a sufficient number of foragers.



Spray paint was either cleaned off or lethal to the bees.



Food coloring works for bees being fed but not the control hives.



Fluorophores have not been tested in honey bees but work very well in other insects.



Chapter 2: Canola


PNW Canola Association



Hypothesis 3: Providing supplemental in-hive feeding will increase the number of foragers working in the field without impacting overall foraging activity.



Year one will utilize a single field.



Hypothesis 4 : Providing supplemental in-hive feeding will result in higher seed purity.




Year two will utilize one field per treatment group



Expected results

3km Test Plot




Year three will combine years one and two



Chapter 3: Colony health



How does feeding during vegetable pollination impact colony health long-term?



Hypothesis 5: Providing supplemental in-hive feeding during vegetable seed pollination will improve colony health long term.





Metrics

- Frames of bees
- Frames of brood
- Frames of food
- Signs of disease
- Varroa mite load



Expected results





Summary

- Movement of pollinators between seed fields decreases seed value
- Honey bees can fly extreme distances when needed
- We want to remove that need
- Bees are really hard to mark effectively



Part 2: Full Hives

Goals

- Increase pollination quality
- Improve colony health
- Updated colony strength recommendations
- Changes to beekeeping practices including the timing of supering



Twohoneys bee co.

Traditionally growers have preferred the hives with the most frames of bees



Image courtesy of popeye.com

Behavioral studies show that colonies forage less under those conditions (Seeley & Tovey, 1994)



Photo by Lily Shui


What we want

What we don't want


Hypothesis: Colonies with more space will exhibit increased foraging activity




Small hive
Small colony




Small hive
Large colony





Large hive
Large colony



How does adding a super impact foraging activity?



Expected results



Summary

- Honey bee foraging behavior decreases as available space in the hive decreases
- The strongest hives may not be the best pollinators
- I will test how adding a super impacts foraging during carrot seed pollination



Thanks



FFAR Fellows
Future Leaders for Food & Agriculture



Questions?

- Riley.reed@wsu.edu
- www.linkedin.com/in/rileymreed
- Feedback survey →
- <https://bugmanriley.com/>



References

- Beekman, M., & Ratnieks, F. L. W. (2000). Long-range foraging by the honey-bee, *Apis mellifera* L. *Functional Ecology*, 14(4), 490–496. <https://doi.org/10.1046/j.1365-2435.2000.00443.x>
- Branchiccela, B., Castelli, L., Corona, M., Diaz-Cetti, S., Invernizzi, C., de la Escalera, G. M., Mendoza, Y., Santos, E., Silva, C., Zunino, P., & Antunez, K. (2019). Impact of nutritional stress on the honeybee colony health. *Scientific Reports*, 9(1), 10156–11. <https://doi.org/10.1038/s41598-019-46453-9>
- Chandel, H., Boring, M., Zentall, T. R., & Wasserman, E. A. (2021). Should I stay or should I go? Pigeons' (*Columba livia*) performance of a foraging task has implications for optimal foraging theory and serial pattern learning. *Journal of Comparative Psychology*, 135(2), 266–272. APA PsycArticles®. <https://doi.org/10.1037/com0000263>
- Couvillon, M. J., Riddell Pearce, F. C., Harris-Jones, E. L., Kuepfer, A. M., Mackenzie-Smith, S. J., Rozario, L. A., Schürch, R., & Ratnieks, F. L. W. (2012). Intra-dance variation among waggle runs and the design of efficient protocols for honey bee dance decoding. *Biology Open*, 1(5), 467–472. <https://doi.org/10.1242/bio.20121099>
- Danner, N., Molitor, A. M., Schiele, S., Härtel, S., & Steffan-Dewenter, I. (2016). Season and landscape composition affect pollen foraging distances and habitat use of honey bees. *Ecological Applications*, 26(6), 1920–1929. <https://doi.org/10.1890/15-1840.1>
- (dataset) USDA National Agricultural Statistics Service. (2019). NASS - Quick Stats. USDA National Agricultural Statistics Service. <https://quickstats.nass.usda.gov/>. [Accessed 18 March 2023].
- Gameda, T. K., Li, J., Luo, S., Yang, H., Jin, T., Huang, J., & Wu, J. (2018). Pollen trapping and sugar syrup feeding of honey bee (Hymenoptera: Apidae) enhance pollen collection of less preferred flowers. *PLOS ONE*, 13(9), e0203648. <https://doi.org/10.1371/journal.pone.0203648>
- Goodwin, R. M. (1986). Increased kiwifruit pollen collection after feeding sugar syrup to honey bees within their hive. *New Zealand Journal of Experimental Agriculture*, 14(1), 57–61. <https://doi.org/10.1080/03015521.1986.10426125>
- Goodwin, R. M., Houten, A. T., & Perry, J. H. (1994). Effect of feeding pollen substitutes to honey bee colonies used for kiwifruit pollination and honey production. *New Zealand Journal of Crop and Horticultural Science*, 22(4), 459–462. <https://doi.org/10.1080/01140671.1994.9513858>
- Goyret, J., & Farina, W. M. (2005). Non-random nectar unloading interactions between foragers and their receivers in the honeybee hive. *Naturwissenschaften*, 92(9), 440–443. <https://doi.org/10.1007/s00114-005-0016-7>
- Hagler, J. R., Hull, A. M., Casey, M. T., & Machtle, S. A. (2021). Use of a Fluorophore to Tag Arthropods for Mark-Release-Recapture Type Research. *Journal of Insect Science*, 21(6), 20. <https://doi.org/10.1093/jisesa/ieab099>
- Hendriksma, H. P., & Shafir, S. (2016). Honey bee foragers balance colony nutritional deficiencies. *Behavioral Ecology and Sociobiology*, 70(4), 509–517. <https://doi.org/10.1007/s00265-016-2067-5>
- Hines, O. (2021, April 13). *Canola Stewardship Statement April 13 2020*. <https://agriculture.basf.us/content/basf/cxm/agriculture/us/en/agriculture/crop-protection/Campaigns/canola-stewardship/canola-stewardship-media-statements/canola-stewardship-update-1.html>
- Klein, A.-M., Vaissière, B. E., Cane, J. H., Steffan-Dewenter, I., Cunningham, S. A., Kremen, C., & Tscharntke, T. (2007). Importance of pollinators in changing landscapes for world crops. *Proceedings of the Royal Society B: Biological Sciences*, 274(1608), 303–313. <https://doi.org/10.1098/rspb.2006.3721>
- Kohl, K. D., Coogan, S. C. P., & Raubenheimer, D. (2015). Do wild carnivores forage for prey or for nutrients? *BioEssays*, 37(6), 701–709. <https://doi.org/10.1002/bies.201400171>
- Månefjord, H., Müller, L., Meng, L., Salvador, J., Blomqvist, S., Runemark, A., Kirkeby, C., Ignell, R., Bood, J., & Brydegaard. (2022). 3D-Printed Fluorescence Hyperspectral Lidar for Monitoring Tagged Insect Movement. *In Revision*.
- Marden, J. H., & Waddington, K. D. (1981). Floral choices by honeybees in relation to the relative distances to flowers. *Physiological Entomology*, 6(4), 431–435. <https://doi.org/10.1111/j.1365-3032.1981.tb00658.x>
- Rivera, M. D., Donaldson-Matasci, M., & Dornhaus, A. (2015). Quitting time: When do honey bee foragers decide to stop foraging on natural resources? *Frontiers in Ecology and Evolution*, 3. <https://www.frontiersin.org/articles/10.3389/fevo.2015.00050>
- Scofield, H. N., & Mattila, H. R. (2015). Honey Bee Workers That Are Pollen Stressed as Larvae Become Poor Foragers and Waggle Dancers as Adults. *PLoS One*, 10(4), e0121731–e0121731. <https://doi.org/10.1371/journal.pone.0121731>
- Visscher, P. K., & Seeley, T. D. (1982). Foraging strategy of honeybee colonies in a temperate deciduous forest. *Ecology (Durham)*, 63(6), 1790–1801. <https://doi.org/10.2307/1940121>
- Zarchin, S., Dag, A., Salomon, M., Hendriksma, H. P., & Shafir, S. (2017). Honey bees dance faster for pollen that complements colony essential fatty acid deficiency. *Behavioral Ecology and Sociobiology*, 71(12), 172. <https://doi.org/10.1007/s00265-017-2394-1>